

# LACTATE OXIDASE [LOX II]

from *Aerococcus viridans*  
(L-Lactate: oxygen oxidoreductase, EC 1.13.12.4)



## Preparation and Specification


Appearance	: Yellowish amorphous powder, lyophilized
Specific activity	: More than 20 U/mg solid
Contaminants	:
POP	Less than 0.001 % (U/U)
GOD	Less than 0.001 % (U/U)
UODN	Less than 0.001 % (U/U)
CO	Less than 0.001 % (U/U)


## Properties

Molecular weight	: 80 kDa (gel filtration)	
Isoelectric point	: pH 4.6	
Michaelis constant	: L-Lactate $7.0 \times 10^{-4}\text{M}$	Figure 1
Optimum pH	: 6.0–7.0	Figure 2
pH stability	: 6.0–9.0 (50°C, 10 min)	Figure 2
Optimum temperature	: 35°C	
Thermal stability	: Stable at 50°C and below (pH 7.0, 10 min)	Figure 3
Storage stability	: At least one year at -20°C	Figure 4

## Applications for Diagnostic Test

This enzyme is useful for enzymatic determination of **lactic acid**.






## Assay

### Principle

The assay is based on the increase in absorbance at 565 nm as the formation of quinoneimine dye proceeds in the following reactions:



DMA: N, N'-Dimethylaniline

### Unit definition

One unit is defined as the amount of enzyme which generates 1  $\mu\text{mole}$  of  $\text{H}_2\text{O}_2$  per minute at 37°C under the conditions specified in the assay procedure.

### Reagents

- Reaction mixture
 

0.2 M KH <sub>2</sub> PO <sub>4</sub> -NaOH buffer pH 6.5	0.20 ml
50 U/ml POD solution <sup>1)</sup>	0.10 ml
15 mM 4-AA solution	0.10 ml
0.5 M DL-Lactic acid solution pH 6.5	0.10 ml
Distilled water	0.30 ml

 Mix above reagents in advance. Just before measuring, add the reagent listed below and mix.
 

0.2% (W/V) DMA solution	0.20 ml
-------------------------	---------

 1): 50 U/ml POD solution  
 Dissolve 500 U (PPU) of POD with 10 ml of distilled water.
- Reaction stopper
 

0.25% (W/V) LBS solution	
LBS: Sodium lauryl sulfate	
- Enzyme dilution buffer
 

10 mM KH <sub>2</sub> PO <sub>4</sub> -NaOH buffer pH 7.0 containing	
10 $\mu\text{M}$ FAD	
FAD: Flavine adenine dinucleotide	

### 4. Reagents

- DL-Lactate: Wako Pure Chemical Industries, Ltd.  
 Special grade #128-00056  
 DMA: Wako Pure Chemical Industries, Ltd.  
 Special grade #044-02763  
 FAD (2Na): Kyowa Hakko Co., Ltd.  
 LBS: NACALAI TESQUE, INC. Extra pure #20123-22  
 4-AA: NACALAI TESQUE, INC.  
 Special grade #01907-52  
 POD: Sigma Chemical Co. Type II #P-8250

### Enzyme solution

Accurately weigh about 20 mg of the sample and add enzyme dilution buffer to make a total of 20 ml. Dilute it with enzyme solution buffer to adjust the concentration as required.

### Procedure

- Pipette accurately 1.0 ml of reaction mixture into a small test tube and preincubate at 37°C.
- After 5 min, add exactly 20  $\mu\text{l}$  of enzyme solution and mix to start the reaction at 37°C.  
 ※ In the case of a test blank, add 20  $\mu\text{l}$  of enzyme dilution buffer in place of enzyme solution.
- At 10 min after starting the reaction, add 2.0 ml of the reaction stopper to stop the reaction.
- Measure the absorbance at 565 nm.
 

Absorbance sample	: A <sub>s</sub>
blank	: A <sub>b</sub>

$$\Delta A = (A_s - A_b) \leq 0.350 \text{ Abs}$$

### Calculation

$$\text{Activity (U/mg of powder)} = \frac{\Delta A / 10}{35.33 \times 1/2} \times \frac{3.02}{0.02} \times \frac{1}{X}$$

35.33 : millimolar extinction coefficient of quinonediimine dye at 565 nm ( $\text{cm}^2 / \mu\text{mole}$ )

1/2 : a multiplier derived from the fact that 2 mole of

$H_2O_2$  produces 1 mole of quinonediimine dye

- 10 : reaction time (min)  
 3.02 : final volume (ml)  
 0.02 : volume of enzyme solution (ml)  
 X : concentration of the sample in enzyme solution  
 (mg/ml)

## Storage

Storage at  $-20^{\circ}C$  in the presence of a desiccant is recommended. The enzyme activity will be retained for at least one year under this condition (Figure 4).

## References

1. Eichel, H. J. and Rem, L. T. (1962) J. Biol. Chem., **237**, 940-945.
2. Esders, T. W. and Goodhue, C. T. (1980) Eastman Kodak Company, U. S. Pat. 4,241,178.

## LOX II 活性測定法 (Japanese)

### I. 試薬液

1. 反応試薬混合液
 

0.2M $KH_2PO_4$ -NaOH 緩衝液 pH6.5	0.20 ml
50U/ml POD 溶液 <sup>1)</sup>	0.10 ml
15mM 4-AA 溶液	0.10 ml
0.5M DL-乳酸溶液 pH6.5	0.10 ml
精製水	0.30 ml

を混合して置く。測定直前に前溶液と  
 0.2% (V/V) DMA 溶液 0.20 ml  
 を混合する。  
 1): 50U/ml POD 溶液  
 POD 500 単位 (PPU) を精製水 10ml で溶解する。
2. 反応停止液  
 0.25% (W/V) LBS 溶液
3. 酵素溶解希釈用液  
 10  $\mu$ M FAD を含む 10mM  $KH_2PO_4$ -NaOH 緩衝液 pH7.0
4. 試薬
 

POD: シグマ社製 Type II #P-8250  
 4-AA: ナカライテスク社製 特級 #01907-52  
 乳酸 (DL-Lactate):  
 和光純薬工業製 特級 #128-00056  
 DMA (N,N'-ジメチルアニリン):  
 和光純薬工業製 特級 #044-02763  
 FAD (フラビンアデニンジヌクレオチド $\cdot$ 2Na):  
 協和発酵製  
 LBS (ラウリルベンゼンスルホン酸):  
 ナカライテスク社製 Extra pure #20123-22

### II. 酵素試料液

検品約 20mg を精密に量り、酵素溶解希釈用液で溶解して全容 20ml とする。  
 その液を酵素溶解希釈用液で適宜希釈する。

### III. 測定操作法

1. 小試験管に反応試薬混合液 1.0ml を正確に分注して  $37^{\circ}C$  で予備加温する。
2. 5 分経過後、酵素試料液 20  $\mu$ l を加えて混和し、 $37^{\circ}C$  で反応を開始する。  
 ※盲検は酵素試料液の代わりに酵素溶解希釈用液 20  $\mu$ l を加える。
3. 10 分経過後、反応停止液 2.0ml を加えて混和し、反応を停止する。
4. 565nm における吸光度を測定する  
 求められた吸光度を試料液は  $A_s$ 、盲検液は  $A_b$  とする。

$$\Delta A = (A_s - A_b) \leq 0.350 \text{ Abs}$$

### IV. 計算

$$\text{活性 (U/mg)} = \frac{\Delta A / 10}{35.33 \times 1/2} \times \frac{3.02}{0.02} \times \frac{1}{X}$$

- 35.33: キノンジイミン色素の 565nm におけるミリモル分子吸光数 ( $cm^2 / \mu$ mole)  
 1/2 :  $H_2O_2$  2 モルからキノンジイミン色素 1 モルが生  
 成することによる係数  
 10 : 反応時間 (min)  
 3.02 : 反応総液量 (ml)  
 0.02 : 反応に供した酵素試料液量 (ml)  
 X : 酵素試料液中の検品濃度 (mg/ml)